

****REPRESENTATIVE DATASHEET****

Sheep anti-human Factor XI (FXI)

Whole IgG from antiserum

5 mg

Product #: SAFXI-IG

Lot #: XXXX

Expiry date: XXXX

Store at -10 to -20°C

 1348 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5
 905-304-9896 • 800-903-6020 • fax 905-304-9897

For Research Use Only.

Not for use in diagnostic procedures.

Description of Factor XI (FXI)

Factor XI (FXI, plasma thromboplastin antecedent) is a coagulation protein produced in the liver that circulates in plasma at approximately 5 µg/ml (30 nM). The mass of FXI is 160 kDa as determined by SDS-PAGE under non-reducing conditions and 80 kDa upon reduction. FXI consists of two identical 80 kDa subunits linked by disulphide bonds. Each subunit consists of a tandem repeat of four apple domains followed by a serine protease catalytic domain. Cleavage of FXI by activated factor XII or thrombin converts each subunit into a two-chain form and generates two active sites per FXIa molecule. The mass of FXIa is 160 kDa unreduced, but upon reduction F.XIa migrates as a heavy chain of 50 kDa and a light chain of 30 kDa. The catalytic site of FXIa resides in the light chain. In plasma, FXI or FXIa circulates in non-covalent 1:1 complex with high molecular weight kininogen, which acts as a cofactor in the activation of FXI by activated factor XII. The activity of FXIa is regulated by platelets and by several proteinase inhibitors including, in order of decreasing importance, C1-inhibitor, α₂antiplasmin, α₁antitrypsin and antithrombin. Heparin has relatively little effect on the rate of inhibition of FXIa by antithrombin. The only known natural substrate for activated FXI (FXIa) is factor IX (Christmas factor) and the only cofactor required for this reaction is ionized calcium¹⁻³.

REFERENCES and REVIEW

1. Willemin WA, Minnema M, Meijers JCM, Roem D, Erenberg AJM, Nuijens JH, ten Cate H, Hack EC; Inactivation of Factor XIa in Human Plasma Assessed by Measuring Factor XIa-Protease Inhibitor Complexes: Major Role for C1-Inhibitor. *Blood* 85:1517, 1995.
2. DeLa Cadena R, Watchfogel YT, Colman RW, in Hemostasis and Thrombosis, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 219-240, J.B. Lippincott Co., Philadelphia, 1994.
3. Baglia FA, Seaman FS, Walsh, PN; The Apple 1 and 4 domains of Factor XI Act to Synergistically Promote the Surface-Mediated Activation of Factor XI by Factor XIIa. *Blood* 85:2078, 1995.

Product Specifications

Description:

Vial containing XXXX ml of whole IgG. Total protein is 5 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Sheep

Immunogen:

Human Factor XI purified from plasma.

Concentration:

 IgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient (E^{1%₂₈₀}) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for FXI as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of antibodies to human FXI.

Neutralizing activity:

XXXX Bethesda Units/ml IgG against normal plasma.

 One Bethesda unit/ml is defined as the amount of inhibitor that resulted in 50% residual FXI activity after 2 hours at 37°C (Kasper CK *et al*, *Thromb Diath Haemorrh* 34:869, 1975).

Species Cross Reactivity: (immunodiffusion vs. citrated plasma)

Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX

Related Products:

Cat #:	GAFXI-IG	Goat anti-human FXI, IgG from antiserum
Cat #:	GAFXI-AP	Goat anti-human FXI, affinity purified IgG
Cat #:	GAFXI-HRP	Goat anti-human FXI, IgG-peroxidase conjugate
Cat #:	SAFXI-AP	Sheep anti-human FXI, affinity purified IgG
Cat #:	FXI-EIA	Paired antibody set for ELISA of FXI, 5 x 96 wells
Cat #:	FXI-DP	Human plasma deficient in FXI, immune depleted
Cat #:	FXI-AG	VisuLize FXI Antigen ELISA Kit

 Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:
AFFINITY BIOLOGICALS INC.
 1348 Sandhill Drive
 Ancaster ON CANADA L9G 4V5
 Tel: (905) 304-9896
 (800) 903-6020
 Fax: (905) 304-9897
info@affinitybiologicals.com