

****REPRESENTATIVE DATASHEET**
Sheep anti-human Protein S (PS)

 Whole IgG from antiserum
 10 mg

Product #: SAPS-IG
Lot #: XXXX
Expiry date: XXXX

Store at -10 to -20°C

 1348 Sandhill Drive, Ancaster, Ontario, Canada L9G 4V5
 905-304-9896 • 800-903-6020 • fax 905-304-9897

 For Research Use Only.
 Not for use in diagnostic procedures.

Description of Protein S (PS)

Protein S (PS) is a vitamin K-dependent glycoprotein produced in the liver, endothelium and megakaryocytes. The concentration of PS in plasma is ~25 µg/ml (~325 nM) where it acts as a cofactor in the anticoagulant activity of activated Protein C. A deficiency of Protein S (quantitative or qualitative) is a risk factor for vascular thrombosis. Protein S is expressed as a single chain molecule with a molecular weight of 77 kDa. The structure of PS is similar to many other vitamin-K dependent coagulation proteins, consisting of an N-terminal calcium binding domain of 10 γ-carboxyglutamic acid (gla) residues, followed by a thrombin-sensitive loop region and 4 EGF-like domains. The C-terminal domain does not contain the usual catalytic triad of a proenzyme, but seems instead to be involved in the binding of PS to C4b-binding protein (C4bp). Protein S binds to activated Protein C (APC) in the presence of calcium and negatively charged phospholipid surface to allow APC to proteolytically inactivate coagulation cofactors Va and VIIIa. Enzymatic regulation of PS cofactor activity is through cleavage of PS in the thrombin-sensitive loop region by thrombin or other enzymes, resulting in the loss of calcium binding properties and APC cofactor activity. Another regulatory mechanism is to reduce the availability of PS by the binding of PS to C4bp. In plasma, approximately 60% of Protein S circulates in non-covalent complex with C4bp, making it unavailable for APC cofactor activity. The binding of PS to C4bp may be important in localizing C4bp to damaged cell membranes where it may control activation of complement by the classical pathway¹⁻³.

REFERENCES and REVIEWS

1. Broze GJ, Miletich JP; Biochemistry and Physiology of Protein C, Protein S and Thrombomodulin; in Hemostasis and Thrombosis, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp 259-276, J.B. Lippincott Co., Philadelphia PA, USA, 1994.
2. Comp PC, Doray D, Patton D, Esmon CT; An Abnormal Plasma Distribution of Protein S Occurs in Functional Protein S Deficiency. Blood 67, pp 504-508, 1986.
3. Schwalbe RA, Dahlback B, Nelsestuen GL; Independent Association of Serum Amyloid P Component, Protein S and Complement C4b with C4b-binding Protein and Subsequent Association of the Complex with Membranes; JBC 265, pp 21749-21757, 1990.

Product Specifications
Description:

Vial containing XXXX ml of whole IgG representing approximately 1 ml of antiserum. Total protein is 10 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Sheep

Immunogen:

Human Protein S purified from plasma.

Concentration:

 IgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ($E^{1\%}_{280}$) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for Protein S as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of antibodies to human Protein S.

Neutralizing activity:

Not determined.

Species Cross Reactivity: (immunodiffusion vs. citrated plasma)

Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX

Related Products:

Cat #: **GAPS-IG** Goat anti-human Protein S, IgG from antiserum
 Cat #: **GAPS-AP** Goat anti-human Protein S, affinity purified IgG
 Cat #: **GAPS-HRP** Goat anti-human Protein S, peroxidase labelled IgG
 Cat #: **SAPS-AP** Sheep anti-human Protein S, affinity purified IgG
 Cat #: **SAPS-HRP** Sheep anti-human Protein S, peroxidase labelled IgG
 Cat #: **PS-EIA** Paired antibody set for ELISA of Protein S, 5 x 96 wells
 Cat #: **PS-DP** Human plasma deficient in PS, immune depleted
 Cat #: **PS-LDP** Lyophilized human plasma deficient in PS, immune depleted

 Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:
AFFINITY BIOLOGICALS INC.
 1348 Sandhill Drive
 Ancaster ON CANADA L9G 4V5
 Tel: (905) 304-9896
 (800) 903-6020
 Fax: (905) 304-9897
info@affinitybiologicals.com
 Document OPI0123, rev 2