

**Catalogue No.**

**Qty:**

600 µg

## Anti-Rab10

**Source:** Goat

**General description:** RAB10 belongs to the large RAB family of low molecular weight GTPases that are involved in intracellular membrane trafficking. Rab10 might be involved in vesicular trafficking and neurotransmitter release.

**Alternative names:** N/A

**Form:** Polyclonal antibody supplied as a 200 µl (3 mg/ml) aliquot in PBS, 20% glycerol and 0.05% sodium azide. This antibody is epitope-affinity purified from goat antiserum.

**Immunogen:** Purified recombinant peptide derived from within residues 100 aa to the C-terminus of mouse Rab10 produced in E. coli.

**Specificity:** Detects Rab10 by Western blot in transfected cells with GFP-Rab10.

**Reactivity:** Reacts with Human, Rat, Mouse, Monkey and Canine proteins

Sample	WB	IHC (F)	IHC (P)	IF	ELISA
Human	+++	ND	ND	ND	ND
Rat	+++	ND	ND	ND	ND
Mouse	+++	ND	ND	ND	ND
Canine	+++	ND	ND	ND	ND
Monkey	+++	ND	ND	ND	ND


+++ excellent, ++ good, + poor, ND not determined

**Usage:**

WB: 1:250-1:2,000

**Storage:** For continuous use, store at 2-8 C for one-two days. For extended storage, store in -20 C freezer. Working dilution samples should be discarded if not used within 12 hours.

**Special instructions:** The antibody solution should be gently mixed before use..



Anti-Rab10 Ab at 1/500 dilution; 293HEK transfected with GFP-Rab10; lysates at 100 µg per lane; rabbit polyclonal to goat IgG (HRP) at 1/10,000 dilution;

For research use only, not for diagnostic use

**SICGEN's Proprietary Immunogen Policy**

In order to produce high specific antibodies SICGEN has invested a lot of time and effort into selecting immunogen sequences. SICGEN has decided to protect this information by not publishing it on the website. However, these sequences are available on request.