

FITC goat anti-rabbit IgG (H+L) *Cross Adsorbed*

Catalog number: 16876

Unit size: 1 mg

Product Details

Storage Conditions 2-6°C and kept from light. To extend the shelf-life of this product, add

an equal volume of glycerol to make a final concentration of

approximately 50% glycerol and store at -20°C.

Expiration Date 12 months upon receiving

Concentration 1 mg/mL

Formulation PBS, 2 mg/mL BSA

Unit Details

Unit 16876 (1 mg)

Reconstitution Volume 1 mL ddH₂O

Antibody Properties

Species Reactivity Rabbit

Class Secondary

Clonality Polyclonal

Host Goat

Chemical Properties

Molecular Weight ~150000

Biological Properties

Stabilizer None

Preparation Goat anti-rabbit IgG (H+L) is produced in goat with pooled total rabbit

IgG, and affinity purified with rabbit IgG coupled beads. The purified IgG has a minimal cross-reaction to human, horse, mouse and bovine IgG.

The antibody is conjugated with FITC under optimal condition. $\label{eq:fitting} % \begin{subarray}{ll} \end{subarray} \begin{subarray}{$

Application Immunofluorescence (IF), Flow Cytometry (FACS)

Soluble In Water

Spectral Properties

Conjugate FITC

Excitation Wavelength 491 nm

Emission Wavelength 516 nm

Applications

AAT Bioquest's anti-mouse secondary antibodies are affinity-purified antibodies with well-characterized specificity for mouse immunoglobulins and are useful in the detection, sorting or purification of its specified target. This FITC-labeled secondary antibody was prepared using AAT Bioquest's proprietary labeling technology. It demonstrated much brighter signal compared to the similar FITC goat anti-mouse IgG antibodies from other commercial sources, thus can significantly increase assay sensitivities. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies can bind to a single primary antibody.