

## iFluor™ 820 goat anti-rabbit IgG (H+L) \*Cross Adsorbed\*

Catalog number: 48060, 48061 Unit size: 200 µg, 1 mg

Product Details			
Storage Conditions	2-6°C and kept from light. To extend the shelf-life of this product, add an equal volume of glycerol to make a final concentration of approximately 50% glycerol and store at -20°C.		
Expiration Date	12 months upon receiving		
Concentration	1 mg/mL		
Formulation	PBS, 2 mg/mL BSA		
Unit Details			
Unit	48060 (200 μg)	48061 (1 mg)	
Reconstitution Volume	$200 \ \mu L \ ddH_2O$	1 mL ddH <sub>2</sub> O	
Antibody Properties			
Species Reactivity	Rabbit		
Class	Secondary		
Clonality	Polyclonal		
Host	Goat		
<b>Chemical Properties</b>			
Molecular Weight	~150000		
<b>Biological Properties</b>			
Stabilizer	None		
Appearance	Green solid		
Preparation	Goat anti-rabbit IgG (H+L) is produced in goat with pooled total rabbit IgG, and affinity purified with rabbit IgG coupled beads. The purified IgG has a minimal cross-reaction to human, horse, mouse, human and bovine IgG. The antibody is conjugated with iFluor™ 820 under optimal condition.		
Application	Flow Cytometry (FACS), ELISA, HC, Western Blot		
Soluble In	Water		
Spectral Properties			
Conjugate	iFluor™ 820		

Excitation Wavelength	822 nm
Emission Wavelength	850 nm

## Applications

AAT Bioquest's iFluor<sup>™</sup> dyes are optimized for labeling proteins, in particular, antibodies. These dyes are bright, photostable, and have minimal quenching on proteins. They can be well excited by the major laser lines of fluorescence instruments (e.g., 350, 405, 488, 532-561, 633-647, and 808 nm). iFluor<sup>™</sup> 820 goat anti-rabbit IgG (H+L) conjugate has fluorescence excitation and emission maxima of 🛛 822 nm and 🖾 850 nm, respectively. These unique spectral characteristics makes iFluor<sup>™</sup> 820 goat anti-rabbit IgG (H+L) conjugate har struments (e.g., and the major laser lines of fluorescence excitation and emission maxima of the major laser lines are specified by the major laser lines of fluorescence excitation and emission maxima of the major laser lines are specified by the major laser lines of fluorescence excitation and emission maxima of the major laser lines are specified by the major laser lines of fluorescence excitation and emission maxima of the major laser lines are specified by the major laser lines are specified by the major laser lines of fluorescence excitation and emission maxima of the major laser lines are specified by the major lines