

Catalogue No.

AB200550-100

Qty:

500 µg

Anti-Goat IgG, DyLight®550

Source: Chicken

General description: Chicken polyclonal antibody to goat IgG conjugated to DyLight®550.

Alternative names: Immunoglobulin Y antibody.

Form: Polyclonal antibody supplied as a 200 µl (2.5 mg/ml) aliquot in PBS, 20% glycerol, 0.05% ProClin® and 0.05% sodium azide.

Immunogen: Full goat IgG affinity purified from domestic goat (*Capra hircus*) serum using protein G. Purity >95% by SDS-PAGE.

Specificity: Using serum samples detects goat IgG by Western blot. Reduced cross-reactivity to other species.

Reactivity: Reacts with Goat proteins

Sample	WB	IHC (F)	IHC (P)	IF	ELISA	IEM
Goat	+++	ND	ND	+++	ND	ND

+++ excellent, ++ good, + poor, ND not determined


Usage:

WB: 1:2,000-1:10,000


IF: 1:100-1:1,000

Storage: Store at -20 C for long-term storage. Store at 2-8 C for up to one month.

Special instructions: Avoid freeze/thaw cycles..



Anti-GAPDH Ab at 1/2,500 dilution using HEK293 transfected cell lysates at 50 µg per lane; chicken polyclonal to goat IgG conjugated to DyLight® 550 (AB200550) at 1/10,000 dilution;



Immunofluorescence in Drosophila larvae NMJ muscle 6/7 expressing GluRIIA-GFP in neurons (GluRIIA is a post-synaptic protein) using 1st Ab anti-GFP at 1/1,000 and 2nd Ab anti-goat IgY conjugated to DyLight®550 at 1/500;

For research use only, not for diagnostic use

SICGEN's Proprietary Immunogen Policy

In order to produce high specific antibodies SICGEN has invested a lot of time and effort into selecting immunogen sequences. SICGEN has decided to protect this information by not publishing it on the website. However, these sequences are available on request.