

**Catalogue No.**

AB6119-100

**Qty:**

300 µg

## Anti-Aquamarine

**Source:** Goat

**General description:** Goat polyclonal antibody to Aquamarine (Aquamarine fluorescent protein). Aquamarine is a basic (constitutively fluorescent), monomeric engineered derivative of green fluorescent protein (GFP) isolated from *Aequorea victoria*. Aquamarine is a ~27 kDa protein that is optimally excited at a 430 nm and has a maximum of emission at 474 nm. It is used in research as a reporter to label and study the biology of the cell using a wide range of applications.

**Alternative names:** green fluorescent protein antibody.

**Form:** Polyclonal antibody supplied as a 100 µl (3 mg/ml) aliquot in PBS, 20% glycerol and 0.05% sodium azide. This antibody is epitope-affinity purified from goat antiserum.

**Immunogen:** Purified recombinant fluorescent protein produced in *E. coli*.

**Specificity:** In 293HEK cells transfected with cds plasmid detects a band of 27 kDa by Western blot. This antibody does not recognize RFP (red fluorescent protein).

**Reactivity:** Reacts with Transfected cells proteins

Sample	WB	IHC (F)	IHC (P)	IF	ELISA	IEM
Transfected cells	+++	+++	+++	+++	ND	+++


+++ excellent, ++ good, + poor, ND not determined

**Usage:**


WB: 1:500-1:5,000  
 IHC (F): 1:50-1:500  
 IHC (P): 1:50-1:500  
 IF: 1:50-1:500  
 IEM: 1:50-1:500

**Storage:** For continuous use, store at 2-8 C for one-two days. For extended storage, store in -20 C freezer. Working dilution samples should be discarded if not used within 12 hours.


**Special instructions:** The antibody solution should be gently mixed before use..



Immunofluorescence – anti-Aquamarine Ab using hCEC cells transduced with Aquamarine-CAV1; cells were fixed with methanol and anti-Aquamarine at 1/250;



Anti-Aquamarine Ab at 1/2,500 dilution using HEK293 transfected cell lysates at 50 µg per lane; rabbit polyclonal to goat IgG (HRP) at 1/10,000 dilution;



Immunofluorescence – anti-Aquamarine Ab using hCEC cells transduced with Aquamarine-CAV1; cells were fixed with methanol and anti-Aquamarine at 1/250;

For research use only, not for diagnostic use

#### SICGEN's Proprietary Immunogen Policy

In order to produce high specific antibodies SICGEN has invested a lot of time and effort into selecting immunogen sequences. SICGEN has decided to protect this information by not publishing it on the website. However, these sequences are available on request.