



Murine Anti-Factor VIII

Clone GMA-8010

Factor VIII (FVIII) is a heterodimer consisting of a heavy chain (ranging in mass from 90 to 200 kDa) bound via metal ions to a light chain (80 kDa). In plasma, FVIII circulates in an inactive form bound to von Willebrand factor. Following activation by factor Xa or thrombin, factor VIIIa can function as cofactor for the enzyme factor IXa in the activation of factor X in the presence of phospholipid and Ca^{2+} . Absent or defective FVIII is the cause of the X-linked recessive bleeding disorder hemophilia A. GMA-8010 recognizes the light chain of FVIII. The antibody inhibits FVIII activation and is suitable for ELISA and bio-layer interferometry pairing experiments.

Description


Antibody Source:	mouse monoclonal, IgG _{2a}
Antigen Species Bound:	human, porcine
Specificity:	FVIII light chain
Immunogen:	B-domain deleted recombinant human FVIII

Formulation and Storage

Purity:	Purified by protein G affinity chromatography from serum-free cell culture supernatant.
Product Formulation:	Lyophilized from a ≥ 1 mg/ml solution in 20 mM NaH_2PO_4 0.15 M NaCl, 1.0% (w/v) mannitol, pH 7.4. Concentration determined by absorbance measurement at 280 nm and using an extinction coefficient of 1.4 ($\epsilon_{0.1\%}$).
Reconstitution:	Reconstitute with deionized water.
Storage:	Store lyophilized or reconstituted and aliquoted material at $-20^\circ C$ for prolonged periods. Avoid freeze-thaw cycles. Alternatively, add 0.02% (w/v) sodium azide to reconstituted solution and store at $4^\circ C$.
Country of Origin:	USA
Size Options:	0.1 mg or 0.5 mg

Applications

Working Concentration:	Approximately 1-5 $\mu g/ml$. Researcher should titer antibody in specific assay.
ELISA:	Binds immobilized human and porcine FVIII.
Immunoblotting:	Not recommended.
Inhibition:	Inhibitory in aPTT clotting assay.
Bio-layer Interferometry:	Can be used in conjunction with GMA-8020, -8012, -8016, -8008, -8013 and -8018 for detection of FVIII.



References

- [1] J.M. Stewart, A.F. Tarantal, W.J. Hawthorne, E.J. Salvaris, P.J. O'Connell, M.B. Nottle, A.J.F. d'Apice, P.J. Cowan, M. Kearns-Jonker. Rhesus monkeys and baboons develop FVIII inhibitors in response to porcine endothelial cells or islets. (2014). *Xenotransplantation*. 21(4):341-352.